

The structure that appears as the large swelling posteriorly on the superior ramus of the ischium and bears most of the weight in sitting, is known as the: - Correct Answer- **Ischial tuberosity**

When percussing the posterior chest, which one of the following techniques would be omitted? - Correct Answer- **Percuss areas over the scapulae**

On examination of the pupils, both are round but the right pupil appears larger

than the left and reacts much slower to light. This condition may be indicative

of: - Correct Answer- **A tonic pupil**

During the type of seizure activity, the patient experiences partial seizures that

resemble tonic-clonic seizures. The patient may recall the aura and unilateral

neurologic deficit is present during the postictal period. This type of seizure activity is referred to as a: - Correct Answer- **focal seizure that becomes generalized**

A female patient complains of weakness in both arms when transferring the

wet clothes from the washer and placing them in the dryer. This finding could

be suggestive of which type of weakness pattern? - Correct Answer-**Symmetric**

To palpate the right ovary when performing the bimanual vaginal exam, the abdominal hand should be on the right lower quadrant and the pelvic hand needs to be: - Correct Answer-**In the right lateral fornix**

A serious disturbance in a person's mental abilities that results in a decreased

awareness of one's environment and confused thinking is referred to as: - Correct Answer-**Delirium**

Which of the following screening tests for hearing loss can detect both sensorineural and conductive hearing loss? - Correct Answer-**Weber test**

Located on the anterior aspect of the distal femur, the patella slides on this groove during flexion and extension of the knee. The name of this groove is the: - Correct Answer-Trochlear groove

A 86-year-old man with uncontrolled hypertension presents with sudden,

intense left lower abdominal pain that radiates to the back. The pain is associated with a tearing sensation. These findings are MOST likely associated

with: - Correct Answer-**An abdominal aortic aneurysm (AAA)**

Ophthalmoscopic examination reveals dark specks noted between the fundus

and the lens. These specks are most likely: - Correct Answer-**Vitreous floaters**

When performing the first Leopold maneuver on a pregnant woman, if the buttocks and head are not easily palpated at the fundus, the fetus is said to be

in: - Correct Answer-**Transverse lie**

When performing a musculoskeletal examination, the nurse practitioner instructs the patient to move his arm in front of his body. This motion of the shoulder girdle would be an example of: - Correct Answer-**Flexion**

To locate the twelfth rib, palpate: - Correct Answer-**Between the spine and the lateral chest**