

**APEA 3P EXAM 1 2025 SPRING –SUMMER
VERIFIED QUESTIONS AND ANSWERS GRADED
A+**

The groove of the metacarpophalangeal joint can be palpated by having the patient...

Flex their hand/spread their fingers

When auscultating breath sounds in a patient who has left sided heart failure, the breath sounds are...

Vesicular with late inspiratory crackles in the dependent portions of the lungs and resonant on percussion

The ankle-brachial index is a screening test used to assess a person's risk for...
Peripheral Artery Disease (PAD)

Olecranon bursitis maybe caused by all of the following except...

Frozen shoulder

In patients who have allergic rhinitis, the nasal mucosa appears...

pale

A term used to describe an increase in muscular bulk with diminished strength is...

Pseudohypertrophy

If abdominal pain persists when the patient raises his head and shoulders, the origin of the tenderness is probably...

In the abdominal wall

On auscultation of the abdomen, rushes of high-pitched sounds are audible and coincide with abdominal cramps. These findings are most consistent with...

Intestinal obstruction

Ophthalmoscopic examination of the fundus reveals tiny, round, red spots in and around the macular area. These findings are consistent with...

microaneurysms

Symptoms of subdural hematoma include...

Noticeable bleeding between the dura and cerebrum on x-ray

On examination of the adult patient, symptoms of fixed posture, tremor, rigidity, and shuffling gait are observed. These findings are consistent with...

Parkinson's Disease

When discussing the musculoskeletal system, all of the following statements related to articular structure disease are true except which one?

Articular disease is usually due to stiffness or pain

Focal tenderness over the trochanter confirms...

Bursitis

Assessing the neurological status of a child with a ventriculoperitoneal shunt should include...

Use of the Glasgow coma scale

The easiest recognizable clinical manifestation(s) of cystic fibrosis in an infant is...

Salty taste on the skin

A reddish blue, irregularly shaped, solid, and spongy mass of blood vessels that may be present at birth and enlarge during the first 10 to 15 months is characteristic of a...

Cavernous hemangioma

The hamstring muscles flex at the knee and are located on the...

Posterior aspect of the thigh

Ophthalmoscopic examination reveals dark specks noted between the fundus and the lens. These specks are most likely...

vitreous floaters